

Branes in the bulk of Anti-de Sitter space

Michael R. Douglas¹ and Washington Taylor IV^{2,3}

¹*Department of Physics and Astronomy
 Rutgers University
 Piscataway, New Jersey 08555, U.S.A.*

and

I.H.E.S., Le Bois-Marie, Bures-sur-Yvette, 91440 France
 mrd@physics.rutgers.edu

²*Department of Physics
 Joseph Henry Laboratories
 Princeton University
 Princeton, New Jersey 08544, U.S.A.*
 wati@princeton.edu

Abstract

We consider $(3 + 1)$ -dimensional $\mathcal{N} = 4$ super Yang-Mills theory with a non-vanishing scalar Higgs vacuum expectation value, and compare this theory to AdS supergravity with branes in the bulk. We show that the one-loop effective potential for excitations of the Yang-Mills field agrees with the classical linearized potential for brane waves in the AdS picture in the limit of long wavelengths. This supports the idea that the AdS/CFT correspondence fits into string theory as expected from previous work.

July 1998

³Address after August 1998: Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge MA 02139, wati@mit.edu

In [1], it was conjectured by Maldacena that there is a remarkable equivalence between $\mathcal{N} = 4$ super Yang-Mills (SYM) theory in $(3 + 1)$ dimensions and type IIB string theory on $AdS_5 \times S^5$. This conjecture has been sharpened and tested in many ways. In the approaches of Gubser, Klebanov and Polyakov [2] and of Witten [3] the connection between SYM theory and AdS string theory is made precise by placing the 3-branes of the SYM theory on the boundary of the AdS space, and associating correlation functions of operators in the boundary SYM theory with supergravity correlation functions subject to certain boundary constraints. In this picture, it is assumed that the adjoint Higgs fields in the SYM theory vanish so that the theory is at a conformal point.

One way to motivate the conjecture is to consider a system of a large number N of coincident D-branes in the string theory. On the one hand, at substringy distances from the branes the dynamics of the system is well-described by truncating to the gauge theory on the branes [4]. On the other hand, for $g_s N \gg 1$ the curvature at these distances is small in string units. This suggests that questions which can be formulated in gravitational terms (e.g., *what is the metric?*) are controlled by supergravity with small stringy corrections, as pointed out in [5].

According to this argument, the AdS/SYM correspondence is a particular limit of the dynamics of branes in string theory, and we can consider other configurations of the branes before taking the limit. In particular, we can consider the Coulomb branch of the gauge theory, i.e. nonzero vacuum expectation values for the scalar fields, corresponding to branes in the bulk of the AdS space. This possibility was also suggested in [1] but it has not played a significant part in the developments so far; in fact, the general belief at this point seems to be that the SYM/AdS connection only makes sense when the scalar vevs vanish.

In this letter we exhibit an example in which this possibility is realized – an SYM configuration with nonvanishing scalar expectation values corresponds to AdS supergravity with branes in the bulk.

We will work with large N $(3 + 1)$ -dimensional SYM theory in a supersymmetric vacuum in which the adjoint Higgs fields take the form

$$X_i = \begin{pmatrix} x_i & 0 & 0 \\ 0 & \tilde{x}_i & 0 \\ 0 & 0 & \mathbf{0} \end{pmatrix}$$

Here $\mathbf{0}$ indicates the vev for $N - 2$ eigenvalues of the Higgs fields, while x_i and $\tilde{x}_i$ are single eigenvalues.

Our basic assertion will be that the eigenvalues x_i and $\tilde{x}_i$ are the positions of 3-branes in the usual way, both before and after taking the scaling limit of [1]. The simplest argument for this is “what else can it be?” – assuming the conjecture, the probe 3-branes must exist in the theory, and there is no other candidate description. However let us see to what extent we can find evidence for this statement without assuming the result.

By using the $O(6)$ symmetry of the theory we can rotate the fields into the form

$$X_1 = \begin{pmatrix} r & 0 & 0 \\ 0 & \tilde{r} \cos \theta & 0 \\ 0 & 0 & \mathbf{0} \end{pmatrix} \quad X_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \tilde{r} \sin \theta & 0 \\ 0 & 0 & \mathbf{0} \end{pmatrix} \quad (1)$$

where $r, \tilde{r}$ are the magnitudes of the 6-vectors $x, \tilde{x}$ and θ is the angle between these vectors.

We now consider the fields (1) as a background and add fluctuations of the gauge field on the two branes. Denoting the $U(1)$ field strength on the first (second) brane by $\hat{F}$ ($\tilde{F}$) respectively, the one-loop effective action describing the leading interaction between the two fields is proportional to [6, 7]

$$\int \prod_i d^4 p_i \left[F^\mu{}_\nu(p_1) F^\nu{}_\lambda(p_2) F^\lambda{}_\kappa(p_3) F^\kappa{}_\mu(p_4) - \frac{1}{4} (F^\mu{}_\nu(p_1) F^\nu{}_\mu(p_2)) (F^\kappa{}_\lambda(p_3) F^\lambda{}_\kappa(p_4)) \right] \\ \times \delta^4(p_1 + p_2 + p_3 + p_4) [G(p_1, p_2, p_3, p_4) + \text{permutations}]$$

where $F = \hat{F} - \tilde{F}$, and

$$G(p_1, p_2, p_3, p_4) = \int d^4 k \frac{1}{(\rho^2 + k^2)(\rho^2 + (k - p_1)^2)(\rho^2 + (k - p_1 - p_2)^2)(\rho^2 + (k + p_4)^2)} \quad (2)$$

with $\rho = \sqrt{(x_i - \tilde{x}_i)^2}$. If we have $N = 2$, this one-loop result determines the four-derivative terms in the effective action exactly, by the non-renormalization theorem of [8].

In general, at this order the result is a sum of pairwise interactions between the branes. Although this contribution to the interaction between pairs of branes at $\mathbf{0}$ is infrared divergent, the interaction between the two branes at x_i and $\tilde{x}_i$ is not affected by this. We can make a stronger statement if we consider the limit from the Coulomb branch, i.e. taking the $N - 2$ eigenvalues to zero, because away from zero the non-renormalization theorem still holds. Since this is uniform as we take the limit and there is no remaining dimensional parameter to control corrections, it is very plausible that the result remains true in the limit. We will see evidence for this shortly.

This result determines the leading scattering of “brane waves,” small fluctuations of the massless modes on branes one and two. The essential statement we are making at this point is that such modes exist and are created by operators such as $\hat{F}$ and $\tilde{F}$, and are weakly interacting. This is clearly true in the original brane picture and at small $g_s N$ but one might argue the point in gauge theory at large $g_s N$. A gauge theory justification for our statement is that we take g_s small (even when $g_s N$ is large), while the interactions to the conformal sector of the system are controlled by the masses r and $\tilde{r}$. This is not to say that such interactions are negligible but that they can produce the dynamics of branes in a IIB string theory background.

Given this statement, it is clear that the result is an observable scattering amplitude. We are now interested in comparing it with expectations from supergravity; i.e. we consider the two brane waves as sources of stress-energy (and the other supergravity fields) and compute their interactions from this point of view.

We first consider the zero-momentum limit; i.e. the case where the fluctuations of the gauge field are constant on both branes. In this case, there is an effective potential between the branes of the form

$$\frac{1}{\rho^4} \left[F^\mu{}_\nu F^\nu{}_\lambda F^\lambda{}_\kappa F^\kappa{}_\mu - \frac{1}{4} (F^\mu{}_\nu F^\nu{}_\mu) (F^\kappa{}_\lambda F^\lambda{}_\kappa) \right] \quad (3)$$

By explicitly expanding this expression in terms of the field strengths $\hat{F}$ and $\tilde{F}$ on the two branes, and grouping the terms according to their Lorentz index structure, the interaction between the two branes can be expressed in terms of a sum of current-current interactions for the stress tensor as well as the sources for the R-R and other NS-NS fields in the IIB theory. An analogous decomposition was given explicitly in the context of Matrix theory and 11-dimensional supergravity in [9]. In the IIB theory the identification of currents is slightly different from that given in [9], although the essential structure is the same; in this case, all the current-current interactions carry an overall factor of $1/\rho^4$. As a simple example of the types of interactions which appear in this decomposition, there are terms of the form

$$\frac{1}{\rho^4} \left[(\hat{F}^{\mu\nu} \hat{F}_{\nu\mu}) (\tilde{F}^{\lambda\rho} \tilde{F}_{\rho\lambda}) \right] \quad \text{and} \quad \frac{1}{\rho^4} \left[(\hat{F}^{\mu\nu} \hat{F}_{\nu\lambda}) (\tilde{F}^{\lambda\rho} \tilde{F}_{\rho\mu}) \right].$$

These terms correspond to pieces of the effective potential arising from dilaton and graviton exchange [10, 11]. Another example is the term of the form

$$\frac{1}{\rho^4} \left[(\hat{F}^{\mu\nu} \hat{F}_{\nu\lambda} \hat{F}^{\lambda\rho}) (\tilde{F}_{\rho\mu}) \right].$$

As shown in [12, 13], the cubic expression in terms of $\hat{F}$ is the leading operator which couples to the field $B_{\rho\mu}$, for which $\tilde{F}_{\rho\mu}$ acts as the source. Although we are only considering fluctuations of the gauge fields on a pair of single 3-branes, it is straightforward to generalize this discussion to include non-abelian scalar vev's in a pair of 3-brane clusters, which are necessary to describe fields with transverse polarization. For such configurations there are “higher moment” contributions to (3) of the form $F^4 X^n / \rho^{4+n}$, as discussed in [9]. These terms should describe the interactions between the higher partial waves on S^5 discussed, for example, in [10].

For $N = 2$, the SYM theory is just the low energy world-volume dynamics of 3-branes in flat space. The first and second 3-branes are separated by a distance

$$\rho = \sqrt{\sum_i (x_i - \tilde{x}_i)^2} = \sqrt{r^2 + (\tilde{r})^2 - 2r\tilde{r}\cos\theta}.$$

From this point of view, the function $1/\rho^4$ appears because it is the Green's function for the flat-space Laplacian in the six dimensions transverse to the 3-brane:

$$\Delta_{\mathbb{R}^6} = -\frac{1}{r^5} \partial_r r^5 \partial_r - \frac{1}{r^2} \Delta_{S^5}$$

$$\Delta_{\mathbb{R}^6} \frac{1}{\rho^4} \sim \delta^6(x - \tilde{x}).$$

We thus see the usual one-loop agreement between gauge theory and supergravity.

At large N and $g_s N$, and in the scaling limit $r \sim l_s^2 \rightarrow 0$, the AdS interpretation is appropriate. The two probe branes are placed in the $AdS_5 \times S^5$ metric, the scaling limit of the metric around the $N - 2$ branes:

$$ds^2 = r^2(-dt^2 + dx_{\parallel}^2) + \frac{dr^2}{r^2} + d\Omega_5^2. \quad (4)$$

The leading long-distance gravitational interaction between a pair of 3-branes with excited gauge or transverse fields is described (in an appropriate gauge) by a propagator $\mathcal{P}(x; \tilde{x})$ which is the Green's function for the Laplacian in the metric (4). This is a priori different from the flat space prediction.

In the case where the excitations on the 3-branes are constant in the parallel directions $x_{\parallel}$, the propagator must satisfy

$$\frac{1}{\sqrt{-g}} \partial_{\mu} \sqrt{-g} g^{\mu\nu} \partial_{\nu} \mathcal{P}(r, \omega; \tilde{r}, \tilde{\omega}) \sim \delta^6(r - \tilde{r}, \omega - \tilde{\omega}) \quad (5)$$

where $(r, \omega) \in \mathbb{R} \times S^5$. In fact, the flat-space Green's function $1/\rho^4$ turns out to solve (5). This can be seen most directly by working out the $AdS_5 \times S^5$ Laplacian. On functions depending only on the coordinates (r, ω) this Laplacian acts as

$$\begin{aligned} \Delta_{AdS} &= -\frac{1}{r^3} \partial_r r^5 \partial_r - \Delta_{S^5} \\ &= r^2 \Delta_{\mathbb{R}^6}. \end{aligned}$$

Thus the effective potential (3) agrees with expectations from AdS supergravity as well.

Agreement in both limits suggests that the result will agree at arbitrary $g_s N$. This is also true and is the supergravity counterpart of the nonrenormalization conjecture (the generalization we suggested of [8] to the conformal limit). Before rescaling, the metric around N 3-branes is

$$ds^2 = f(r)^{-1/2}(-dt^2 + dx_{\parallel}^2) + f(r)^{1/2}(dr^2 + r^2 d\Omega_5^2)$$

with

$$f(r) = 1 + \frac{g_s N}{r^4}.$$

One can again check that $\Delta_{general} = f^{-1/2} \Delta_{\mathbb{R}^6}$ allowing the same Green's function to solve this Laplacian as well.

We now briefly discuss the case of non-zero momentum. One can check that turning on non-zero momenta in (2) leads to corrections controlled by the scale l_s , for which we have not found a supergravity interpretation. What is clear from the expression (2) is that the result is analytic in the external momenta at $p = 0$, and has an expansion in p/r , as usual in effective field theory. In other words, a derivative d/dp can be expressed in terms of the

derivative d/dr (and less singular terms as $r \rightarrow 0$). As we discuss below, we interpret this in terms of an uncertainty relation for the position of the brane.

In this note we have observed that the leading interaction between brane waves on 3-branes agrees when computed as a quantum effect in gauge theory, and as an interaction mediated by bulk fields in supergravity, both in a flat background and in the AdS background corresponding to a large number of additional 3-branes.

Besides providing a further example of an agreement between super Yang-Mills theory and supergravity, we consider the result to be significant evidence for the following identification: in a supersymmetric vacuum in the gauge theory, a non-zero eigenvalue for the scalar vev corresponds to the position of an *individual* 3-brane. This is the naive correspondence suggested by the brane picture, but it is worth stating it again in the context of the AdS/CFT correspondence.

Note that this does not necessarily mean that we should consider the $N - 2$ branes in our illustration as located at the point $x = 0$. Conceptually, if we are regarding supergravity as coming out of large N gauge theory dynamics, the simplest picture (as has been described by Maldacena; see also [14]) is not to consider the conformal sector of the theory as localized anywhere. Excitations on the $N - 2$ branes are not free brane waves – once we consider $g_s(N - 2)$ large there is no control over infrared divergences in our computation, and no justification for the result (3).

Another class of vacua mentioned in [1], and motivated from the brane point of view is to split the branes into groups with $N = \sum N_i$ each with a different scalar vev. If we take the scaling limit of these vevs, the natural conjecture is that when all $g_s N_i$ become large, this corresponds to supergravity in the background of a multicenter solution. Again one is not asserting that “the N_i branes are located at these points” because their excitations are not brane waves; however a probe brane could be localized. This example has been studied in [18]; many questions remain. We mention that our result also works in this case.

The identification also gives us a class of local observables in AdS space, the operators which create excitations on the probe branes. An important point however is that the localization of the probe branes appears to be precise only in the zero longitudinal momentum limit. In work such as [17, 4] it was observed that to localize the branes in transverse space, one needed arbitrarily low energies, because the energy of the states responsible for the interactions went as $E \sim r/l_s^2$. This can also be formulated as an uncertainty principle [15]: using $\Delta E \Delta t > 1$ we have $\Delta r \Delta t > l_s^2$ and we need arbitrarily large time to make such a measurement. The scaling limit of [1] takes $r \sim l_s^2 \rightarrow 0$, so the uncertainty principle survives, as observed in [16].

We interpret the corrections to the interaction of brane waves at nonzero momentum (compared to supergravity) as a manifestation of this uncertainty principle. Given an uncertainty Δp around zero in energy-momentum, we saw that the difference between the gauge theory and supergravity potentials V can be approximated by our observations as $\Delta p \, d/dr \, V$. This is what would come from an uncertainty $\Delta r \, d/dr \, V$ with $\Delta r \sim \Delta p \sim \Delta E$

as above. It will be interesting to follow this up and especially to understand if this is a universal constraint on our ability to formulate local observables in the AdS language.

Finally, this paper only discussed the case of gauge theory on $\mathbb{R}^{3,1}$, while the AdS/SYM correspondence can be formulated for more general base spaces, for example $S^3 \times \mathbb{R}$. One may ask: how does one describe a probe 3-brane in this case? Doesn't the lifting of the moduli space described in [3] eliminate the configurations we described here?

The answer is clear once one takes the conformal symmetry of the theory into account. In some cases one can relate the theory on different base spaces by conformal transformations, and map the simple supersymmetric vacua we describe here into solutions on a different base space. More generally, one needs to find solutions which preserve 16 of the superconformal symmetries; we predict that appropriate candidates will exist. All of these are time-dependent solutions, so there is no contradiction with the nonexistence of a moduli space.

Applying conformal transformations allows us to produce new solutions in $\mathbb{R}^{3,1}$ as well. Since the corresponding AdS isometries can map points within the coordinate patch (4) to points outside [19], these describe new 3-branes extending across the asymptotic observer's horizon. These will be studied in future work.

Acknowledgements

This work was carried out at the workshop on Dualities in String Theory at the ITP Santa Barbara, and we would like to thank David Gross, Dan Hone and the entire staff for providing a fantastic environment for doing physics.

The work of MRD is supported in part by DOE grant DE-FG02-96ER40559. The work of WT is supported in part by the National Science Foundation (NSF) under contract PHY96-00258.

We also call the reader's attention to [20], which uses 3-branes to study microscopic aspects of the AdS/SYM correspondence.

References

- [1] J. Maldacena, "The large N limit of superconformal field theories and supergravity," [hep-th/9711200](#).
- [2] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, "Gauge theory correlators from non-critical string theory," [hep-th/9802109](#).
- [3] E. Witten, "Anti-de Sitter space and holography," [hep-th/9802150](#).
- [4] M. R. Douglas, D. Kabat, P. Pouliot, and S. Shenker, "D-Branes and Short Distances in String Theory," *Nucl. Phys.* **B485** (1997) 85, [hep-th/9608024](#).

- [5] M. R. Douglas, J. Polchinski and A. Strominger, *JHEP* **12** (1997) 003, [hep-th/9703031](#).
- [6] S. J. Gates Jr., M. T. Grisaru, M. Rocek and W. Siegel, *SUPERSPACE or One thousand and one lessons in supersymmetry*, (Benjamin/Cummings Publishing Company, Reading, 1987).
- [7] V. Periwal and R. von Unge, “Accelerating D-branes,” [hep-th/9801121](#).
- [8] M. Dine and N. Seiberg, “Comments on higher derivative operators in some SUSY field theories,” [hep-th/9705057](#).
- [9] D. Kabat and W. Taylor, “Linearized supergravity from Matrix theory,” *Phys. Lett. B* **426** (1998) 297-305, [hep-th/9712185](#).
- [10] I. R. Klebanov, *Nucl. Phys. B* **496** (1997) 231, [hep-th/9702076](#).
- [11] S. S. Gubser, I. R. Klebanov and A. A. Tseytlin, *Nucl. Phys. B* **499** (1997) 217, [hep-th/9703040](#).
- [12] S. R. Das and S. P. Trivedi, “Three-brane action and the correspondence between $\mathcal{N} = 4$ Yang-Mills theory and anti-de Sitter space,” [hep-th/9804149](#).
- [13] S. Ferrara, M. A. Lledo and A. Zaffaroni, “Born-Infeld Corrections to D3 brane Action in $AdS_5 \times S_5$ and $N=4$, $d=4$ Primary Superfields,” [hep-th/9805082](#).
- [14] T. Banks, M. R. Douglas, G. Horowitz and E. Martinec, to appear.
- [15] T. Yoneya, *Mod. Phys. Lett. A* **4** (1989) 1587; M. Li and T. Yoneya, *Phys. Rev. Lett.* **78** (1997) 1219, [hep-th/9611072](#).
- [16] M. Li and T. Yoneya, “Short-distance space-time structure and black holes in string theory: a short review of the current status,” [hep-th/9806240](#).
- [17] A. Sen, *Nucl. Phys. B* **475** (1996) 562, [hep-th/9605150](#); T. Banks, M. R. Douglas, N. Seiberg, *Phys. Lett. B* **387** (1996) 278, [hep-th/9605199](#); N. Seiberg, *Phys. Lett. B* **384** (1996) 81, [hep-th/9606017](#).
- [18] J. Minahan and N. Warner, “Quark Potentials in the Higgs Phase of Large N Supersymmetric Yang-Mills Theories,” [hep-th/9805104](#).
- [19] G. Horowitz and H. Ooguri, *Phys. Rev. Lett.* **80** (1998) 4116-4118, [hep-th/9802116](#).
- [20] M. Berkooz, [hep-th/9807230](#).